ISSN (print) 0093-4666 © 2015. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

http://dx.doi.org/10.5248/130.437

Volume 130, pp. 437–443 *April–June 2015*

Two new conidial fungi from Chapada Diamantina, Brazil

Carolina Ribeiro Silva[1], Luís Fernando Pascholati Gusmão[2*], & Rafael F. Castañeda-Ruiz[3]

[1] *Depto. de Micologia, Universidade Federal de Pernambuco, 50670-420, Recife, Brazil*

[2] *Depto. de Ciências Biológicas, Lab. de Micologia, Universidade Estadualde Feira de Santana, Avenida Transnordestina s/n, bairro Novo horizonte, 44036-900, Feira de Santana, Brazil*

[3] *Inst. de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt' (INIFAT), Académico Titular de la Academia de Ciencias de Cuba, Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200*

* Correspondence to: *lgusmao@uefs.br*

Abstract —*Phaeoschizotrichum ramosum* gen. & sp. nov. and *Pyriculariopsis bicolorata* sp. nov., two new asexual fungi collected on leaves of *Calophyllum brasiliense* in the Brazilian semiarid Caatinga region, are described and illustrated. *Phaeoschizotrichum ramosum* is characterized by distinctly branched conidiophores, discrete, terminal, sympodially extended conidiogenous cells, and subulate to narrowly obclavate, 1–4-septate, pale brown conidia. *Pyriculariopsis bicolorata* is distinguished by conidia that are obclavate to subulate, 1-septate, bicolored, and smooth.

Key words — microfungi, taxonomy, leaf litter

Introduction

During an investigation of conidial fungi associated with decaying leaves of *Calophyllum brasiliense* Cambess. (*Clusiaceae*) in a riparian forest of Chapada Diamantina (Piatã, Bahia) in the semiarid Caatinga region of Brazil, two interesting fungi were collected. One was remarkably different from all previously described hyphomycete genera (Seifert et al. 2011) while the other differed from all *Pyriculariopsis* species (Iturriaga et al. 2008, Soares et al. 2011). The two fungi are therefore described here as new.

Materials & methods

During studies in periods September 2013 to July 2014 leaves of *C., brasiliense* were collected in Piatã, Bahia. The material was taken to the laboratory using the methodology by Castañeda-Ruiz (2005). Sample leaves of *C. brasiliense* were placed in plastic bags. In

the laboratory the samples were washed, placed in Petri dish moist chambers, and stored in a polystyrene box with sterile water plus glycerol for 30 days. Slide mounts were prepared in PVL (polyvinyl alcohol, lactic acid, and phenol) and measurements were made at a magnification of ×1000. Microphotographs were obtained with an Olympus microscope (model BX51) equipped with bright field and Nomarski interference optics. The type specimens are deposited in the Herbarium of Universidade Estadual de Feira de Santana, Brazil (HUEFS).

Taxonomy

Phaeoschizotrichum C.R. Silva, Gusmão & R.F. Castañeda, **gen. nov.**

MycoBank MB810712

Differs from *Schizotrichum* by its mostly branched conidiophores, its percurrent and sympodially extended small denticulate and slightly melanized conidiogenous cells, and its pale brown to brown conidia.

Type Species: *Phaeoschizotrichum ramosum* C.R. Silva et al.

Etymology: Greek, *Phaeo-*, meaning dark, referring to the pale brown conidial pigmentation; Latin, *-schizotrichum* referring to the hyphomycete genus *Schizotrichum.*

Colonies hairy, effuse, brown. Mycelium superficial. Conidiophores distinct, single, mostly branched above, septate, brown to dark brown. Conidiogenous cells polyblastic, denticulate, integrated or discrete, enteroblastic percurrent elongated and sympodial extensions, terminal. Conidial secession schizolytic. Conidia solitary, subulate, narrow obclavate to subcylindrical, euseptate, pale brown to brown, smooth or verruculose.

Note: *Phaeoschizotrichum* superficially resembles *Schizotrichum* McAlpine (Ellis 1971, Seifert et al. 2011), but *Schizotrichum* has immersed mycelium and forms erumpent dark brown textura angularis stromata; setae are sometimes present and unbranched, dark brown, and thick-walled while its conidiophores are caespitose, unbranched, and bear sympodial elongated conidiogenous cells and subulate or filiform, smooth, hyaline conidia. *Phaeoschizotrichum* also superficially resembles other cercosporoid fungi with pale or dark pigmented conidia that are known to cause foliar diseases, such as *Passalora* Fr. and *Pseudocercospora* Speg.

Phaeoschizotrichum ramosum C.R. Silva, Gusmão & R.F. Castañeda, **sp. nov.**

MycoBank MB810713 Figs 1, 2

Differs from *Schizotrichum lobeliae* by its superficial mycelium composing an anastomosed network of hyphae with a textura epidermoidea appearance, its mostly branched multi-septate ventricose conidiophores, and its pale brown conidia.

Type: Brazil, Bahia, Piatã, Serra da Tromba, 13°07'S 41°50'W, on dead leaf of *Calophyllum brasiliense*, 19 XI. 2013, coll. C.R. Silva (**Holotype**: HUEFS 210423).

Etymology: Latin, *ramosum,* referring to the branched conidiophores.

Fig. 1. *Phaeoschizotrichum ramosum* (ex HUEFS 210423). A–D. Conidia. E–H. Conidiogenous cells. I–K. Conidiophores and conidiogenous cells.



Fig. 2. *Phaeoschizotrichum ramosum* (ex HUEFS 210423). A–B. Hyphae and conidiophore basal cells. C. Conidiophore and conidiogenous cells. D–I. Branches with fissured outer cell walls and conidiogenous cells.

Colonies on the natural substratum, effuse, hairy, amphigenous, brown. Mycelium superficial, composed of septate, pale brown, 2–4 µm wide, smooth, branched hyphae, forming an anastomosed network of textura epidermoidea appearance. Conidiophores distinct, single, rarely unbranched, mostly branched above, erect, cylindrical, smooth near the base, ventricose, closely septate, with fissured outer cell walls near the apex and branches, with up to 10 enteroblastic percurrent extensions in the main axis and branches, brown below, subhyaline to pale brown above, 100–200 × 5–10 µm. Conidiogenous cells polyblastic, cylindrical-subulate, slightly denticulate, weakly melanized at the conidiogenous loci, subhyaline to pale brown, sympodially elongated after enteroblastic percurrent extensions, smooth sometimes cracked, furrowed or leprous near the percurrent elongation, 15–50 × 4–5 µm. Conidia solitary, subulate, narrowly obclavate, truncate, slightly obscure at the base, rounded or obtuse at the apex, 1–4-septate, mostly 3-septate, pale brown, 25–68 × 2.5–4 µm, smooth, rarely verruculose, dry.

Pyriculariopsis bicolorata C.R. Silva, Gusmão & R.F. Castañeda, **sp. nov.** Fig. 3

MycoBank MB810714

Differs from *Pyriculariopsis* spp. by smaller, 1-septate, bicolored conidia.

Type: Brazil, Bahia, Piatã, Serra da Tromba, 13°07'S 41°50'W, on dead leaf of *Calophyllum brasiliense*, 2 X. 2013, coll. C.R. Silva (**Holotype**: HUEFS 210424).

Etymology: Latin, *bicolorata*, referring to the two-colored conidia.

Colonies on the natural substratum, effuse, hairy, hypophyllous, brown. Mycelium superficial and immersed, composed of septate, branched, brown, 1–2 µm wide, smooth hyphae. Conidiophores distinct, single, develop as lateral branch of hyphae, unbranched, cylindrical, straight to slightly flexuous, 4–7-septate, brown dark at the base becoming paler to the apex, with 2–5 percurrent extensions, 75–175 × 2.5–4 µm, smooth. Conidial secession schizolytic. Conidiogenous cells polyblastic, integrated, mostly terminal, sometimes intercalary, sympodially elongated, slightly denticulate, pale brown, 15–47.5 × 2.5–4 µm, slightly cicatrized at the loci. Conidia obclavate, truncate at the base, obtuse and rounded apex, slightly constricted at the septa, 1-septate, bicolored, brown basal cell, subhyaline apical cell, single, dry, 12–18 × 2.5–4 µm.

Note: A key to eight *Pyriculariopsis* species with conidial illustrations was provided by Iturriaga et al. (2008); subsequently Soares et al. (2011) named *P. calatheae*, which is characterized by cylindrical-fusiform to obclavate, 18–36 × 5–7 µm, 1–2-septate, often with a rostrate apical cell, 3–12 × 2–4 µm, pale brown to subhyaline. None of the nine previously described species of *Pyriculariopsis* is close to or resembles *P. bicolorata*.



Fig. 3. *Pyriculariopsis bicolorata* (ex HUEFS 210424). A–E. Conidia. F. Conidium and conidiogenous cells. G–H. Conidiogenous cells. I. Conidiophore with external hyphae.

Acknowledgments

The authors express their sincere gratitude to Dr. De-Wei Li and Dr. Raghvendra Singh for their critical review of the manuscript. The authors are grateful to PPBio Semiárido and Edital Universal do CNPq for financial support (Proc. 558317/2009-0 and 558317/2009-0). The authors thank the Programa de Pós-Graduação em Biologia de Fungos (PPGBF/ UFPE). C.R. Silva and L.F.P. Gusmão extend their gratitude to CNPq for financial support (Proc. 132415/2013-5). RFCR is grateful to Cuban Ministry of Agriculture and "Programa de Salud Animal y Vegetal", project P131LH003033 for facilities. We acknowledge the assistance provided by Dr. P.M. Kirk and Drs. V. Robert and A. Decock through the IndexFungorum and MycoBank websites. Dr. Lorelei Norvell's editorial and Dr. Shaun Pennycook's nomenclatural reviews are greatly appreciated.

Literature cited

Castañeda Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183, in: Anais do V Congresso Latino Americano de Micologia. Brasilia.

Ellis MB. 1971. Dematiaceous hyphomycetes. Kew, Commonwealth Mycological Institute.

Iturriaga T, Fernandéz R, Castañeda-Ruiz RF, Minter DW, Heredia GA. 2008. A new anamorphic fungus from Venezuela: *Pyriculariopsis formosa*. Mycotaxon 105: 337–342.

Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p. http://dx.doi.org/10.3767/003158511X617435

Soares DJ, Rocha FB, Barreto RW. 2011. *Pyriculariopsis calatheae* sp. nov., a novel anamorphic hyphomycete from the Atlantic forest of Brazil causing leaf spots on *Calathea longifolia*. Mycol. Progress 10: 315–321. http://dx.doi.org/10.1007/s11557-010-0704-3